

Protein Unfolding and the Diffusion Collision Model

Chris Beck^a and Xavier Siemens^b

February 1, 2008

*^aMolecular Modeling Laboratory and ^bInstitute of Cosmology
Department of Physics and Astronomy
Tufts University
Medford, MA 02155*

Abstract

In the diffusion-collision model, the unfolding or backward rates are given by the likelihood of secondary structural cluster dissociation. In this work, we introduce a backward rate calculation modeled from a Kramers-type thermal tunneling through a barrier, which represents the free energy potential well for buried hydrophobic residues. Our results are in good agreement with currently accepted values and the approach suggests a link between the diffusion-collision and folding funnel models of protein folding.

I. Introduction

In the diffusion-collision model (DCM) of protein folding (Bashford et al 1988, Karplus and Weaver, 1976, 1979, 1994) the protein is modeled using a collection of spheres connected by flexible strings. The spheres represent the secondary structural elements such as α -helices or β -sheets, called microdomains, that make up the protein.

The folding process from a completely unfolded protein to the the final native state is accomplished via diffusion through the solvent, collision, and finally coalescence of the microdomains. The state of the protein is defined by the number of pairings (hydrophobic interactions) between the microdomains that are present at a given time t . The rate equations for transitions between these states can be written as

$$\frac{d\mathbf{P}(t)}{dt} = \hat{K}\mathbf{P}(t) \quad (1)$$

where $\mathbf{P}(t)$ is the vector of states and $\hat{K}$ is a matrix containing the transition rates between the different states.

As an example, consider a simple monomeric protein containing two α -helices A and B joined by a connecting string. This gives us a one-pair/two-state system. We will call state 1 the state when the two microdomains are not in hydrophobic contact, and state 2 when they are hydrophobically docked. In this case (1) can easily be written out explicitly

$$\begin{aligned} \dot{P}_1(t) &= -k_{1\rightarrow 2}P_1(t) + k_{2\rightarrow 1}P_2(t) \\ \dot{P}_2(t) &= k_{1\rightarrow 2}P_1(t) - k_{2\rightarrow 1}P_2(t) \end{aligned} \quad (2)$$

or in matrix form as

$$\frac{d}{dt} \begin{pmatrix} P_1(t) \\ P_2(t) \end{pmatrix} = \begin{bmatrix} -k_{1\rightarrow 2} & k_{2\rightarrow 1} \\ k_{1\rightarrow 2} & -k_{2\rightarrow 1} \end{bmatrix} \begin{pmatrix} P_1(t) \\ P_2(t) \end{pmatrix}. \quad (3)$$

A more complicated protein having, say, n microdomains would involve $p = n(n-1)/2$ pairings, 2^p states $P_i(t)$ and a $2^p \times 2^p$ rate matrix $\hat{K}$. In general the calculation of the elements of the rate matrix $\hat{K}$ is somewhat involved. The forward rates are the rates of microdomain coalescence. In the diffusion-collision model the forward rates are calculated assuming the microdomains diffuse through a solvent environment, the space of which is limited by the length of the intervening strings and the van der Waals radii of the microdomains. These microdomains are assumed to be nascently

formed, and their degree of formation is given by a helix-coil transition theory (Zimm and Bragg, 1959, Lifson and Roig, 1961) as in AGADIR (Lacroix et al, 1998, Munoz and Serrano, 1994 a,b,c, 1997) calculation in the case of α -helices, or via a combination of theory (Munoz et al, 1998) and experiment (Dinner et al, 1999) in the case of β -sheets. As the microdomains undergo diffusion, they occasionally collide. When this happens the microdomains coalesce with a probability γ , being held together by hydrophobic interactions in the case of α -helices, or a combination of hydrophobic and hydrogen bond interactions in the case of β -sheets. The coalescence probability γ is given by the likelihood that the microdomain is in α -helical or β -sheet form, the percentage of hydrophobic area, and the likelihood of proper geometrical orientation upon collision.

The forward folding times in the mean first passage time approximation (Weiss, 1967, Weaver, 1979, Szabo et al 1980) are given by

$$\tau_f = \frac{l^2}{D} + \frac{LV(1-\gamma)}{\gamma DA} \quad (4)$$

where V is the diffusion volume available to the microdomain pair, A is the target area for collisions, D is the relative diffusion coefficient, γ is the probability of coalescence upon collision and l and L are geometrical parameters calculated for diffusion in a spherical space. The inverse of the first passage time-scales τ_f are the forward folding rates k_f that are used in the rate matrix $\hat{K}$. In the example given by (2) and (3), k_f is $k_{1 \rightarrow 2}$.

The microdomain pairings can also dissociate. To date, in typical diffusion-collision model calculations, the form of the backward folding, or unfolding times τ_b used for two microdomains A and B comes from the Van't Hoff-Arrhenius law (Van't Hoff, 1884, Arrhenius, 1889) given by

$$\tau_b = \nu^{-1} e^{f A_{AB}/k_B T} \quad (5)$$

where f is the free energy change per unit buried hydrophobic area in the pairing (Chothia, 1974), A_{AB} is the buried area (Lee and Richards, 1971), k_B is Boltzmann's constant, T is the temperature and ν is an attempt rate.

In diffusion collision model calculations to date the attempt rate is a parameter whose value must be set by hand, usually requiring some guess work or adjustment to obtain the desired result (see for example Burton et. al 1998). The typical values used lie in the range $1 \times 10^9 s^{-1} - 1000 \times 10^9 s^{-1}$, obtained from estimates of covalent bond oscillation frequencies (Fersht, 1999). This procedure is not explicitly stated in most diffusion-collision model studies; it is justified, however, when the equilibrium occupation probabilities of the states are known. In fact, the rate constants can be calculated from such probabilities (Chandler, 1987). In cases where the final occupation probabilities are unknown, for instance in studies of protein mis-folding and non-native kinetic intermediates (Beck et al., 2000), such methods are clearly not possible.

In this work we show how the parameter ν , or more generally the unfolding rates, can be determined from thermal fluctuations providing a means of avoiding the guesswork. This makes the diffusion-collision model more predictive and enables it to be used in situations where the final occupation probabilities are unknown.

II. Calculation of the Unfolding Rates

In order to calculate the backward rate it is convenient to view the unfolding process as that of diffusion within, and escape from, an effective one-dimensional potential well. This is a good approximation if only motion perpendicular to the hydrophobic contact surface is important in microdomain pair dissociation.

Consider the pair of microdomains A and B connected by a string. Microdomain coalescence and dissociation can be approximated by diffusion in a potential like the one depicted in Figure 1. The quantity that diffuses is the separation between the microdomains and we can think of

$$dP(x) = \rho(x)dx \tag{6}$$

as the fraction of microdomains with a distance between them in the range




Figure 1: Diffusion potential for the two microdomains A and B. The potential is infinite on the far right because the microdomain separation cannot exceed the length of their connecting string. On the left it is also infinite, in this case because of the hard-core repulsion of the van der Waals contact between the microdomains. The potential energy barrier has a height $E_b = fA_{AB}$, the free energy difference between paired and unpaired states with a buried hydrophobic area A_{AB} , and width L , which we take to be the diameter of a water molecule.

x and $x + dx$. In a potential $V(x)$ the probability density $\rho(x)$ satisfies the one dimensional Smoluchowski equation

$$\frac{\partial \rho(x, t)}{\partial t} = \frac{\partial}{\partial x} \left[D \left(\frac{\partial \rho(x, t)}{\partial x} + \rho(x, t) \frac{\partial V(x)}{\partial x} \right) \right]. \quad (7)$$

The outer boundary on the far right is given by an infinite potential, arising from the two microdomains coming to the end of their connecting string. The inner boundary is also infinite, stemming from the hard-core repulsion of the van der Waals contact between the microdomains. The well depth $E_b = fA_{AB}$ is the free energy difference between paired and unpaired microdomains and the well width L is set to the diameter of a water molecule. A separation larger than L exposes the microdomain to the solvent, the free

energy savings is lost, and the microdomains separate, resulting in an escape from the potential well.

The forward folding rates can be satisfactorily found by the mean first passage time using the method outlined above (4). As a consequence of this we shall forego direct analysis of the entire diffusion process and concentrate on the escape of a bound microdomain out of the potential well in order to find the backward folding rate.

If we concentrate on the part of the potential where $x < L$ in Figure 1, we can think of the potential as being perfectly reflecting on one side and partially permeable on the other. The permeability arises from the thermal fluctuations in the energy which allow the microdomains to escape the bounded region, provided they have sufficient energy, and it can be expressed using an appropriate choice of boundary conditions. We proceed as follows.

For the inner boundary, representing van der Waals contact, we impose perfectly reflecting boundary conditions, meaning the flux through the boundary is zero

$$J(0, t) = D \left. \frac{d\rho(x, t)}{dx} \right|_{x=0} = 0. \quad (8)$$

The flux at the permeable boundary at $x = L$ depends on the density of microdomains at that boundary and the probability that the microdomain energy is high enough to thermally tunnel through the boundary. The differential element of flux at the boundary L of microdomains with relative velocity between v and $v + dv$ is given by

$$dJ(L, t) = v\rho(L, t)dN(v) \quad (9)$$

where

$$dN(v) = \left(\frac{\mu}{2\pi k_B T} \right)^{1/2} e^{-\mu v^2 / 2k_B T} dv \quad (10)$$

is the Maxwell-Boltzmann distribution, namely the fraction of microdomains

with relative velocities between v and $v + dv$ and μ is the reduced mass given by

$$\mu = \frac{m_A m_B}{(m_A + m_B)} \quad (11)$$

where m_A and m_B are the masses of the two microdomains. The distribution is normalised so that the bounds of the integral go from $-\infty$ to ∞ .

It is clear that in order to find the total flux through the outer boundary at L we must integrate over all microdomain velocities larger than $+\sqrt{E_b/2m}$ since the potential well height is E_b , and only microdomains with velocity higher than that can escape and thus contribute to the flux leaving the well. This yields a flux out of state 2

$$J_2^{out}(L, t) = \rho(L, t) \left(\frac{k_b T}{2\pi\mu} \right)^{1/2} e^{-E_b/k_B T} \quad (12)$$

If we assume that the probability is uniformly distributed across the well, then $\rho(x, t) = P_2(t)/L$ everywhere within the well. This is a reasonable assumption because the permeability of the well is so low. Furthermore it can be shown that typically the damping time (or velocity autocorrelation) time is considerably smaller than the well crossing time and the mean first passage time. Clearly then, diffusion in the interior of the well distributes the microdomain separations evenly among the available space and therefore

$$J_2^{out}(L, t) = \frac{P_2(t)}{L} \left(\frac{k_b T}{2\pi\mu} \right)^{1/2} e^{-E_b/k_B T}. \quad (13)$$

The interesting result is that by using a stationary flux method (Kramers, 1940), or equivalently setting $P_2(t) = 1$, the outward flux at the boundary in one dimension $J^{out}(L, t)$ is the fraction of microdomain pairs crossing the boundary at L per unit time and thus clearly identical to the unfolding rate k_b for the AB microdomain pairing. Below we explain this in more detail and show more generally how the rate equations of the diffusion-collision model (1, 2) can be derived from the diffusion equation.

To understand this, we remember that the Smoluchowski equation (7) is

simply a detailed statement of the equation of continuity

$$\frac{\partial \rho(\mathbf{x}, t)}{\partial t} = \nabla \cdot \mathbf{J}(\mathbf{x}, t) \quad (14)$$

The quantity in brackets in (7) is the flux $\mathbf{J}(\mathbf{x}, t)$, which includes the ordinary $\nabla \rho(\mathbf{x}, t)$ term, and the second term which takes into account external forces. Integration of both sides of (14) over the bound part of diffusion volume, using the divergence theorem on the R.H.S. and assuming isotropic flow into and out of the boundary of the diffusion volume yields

$$\frac{dP_2(t)}{dt} = \mathbf{n} \cdot \mathbf{J}(t)A = \{J_2^{in}(t) - J_2^{out}(t)\}A. \quad (15)$$

The LHS of this equation is the rate of change of probability in the bound region. On the RHS $\mathbf{n}$ is a unit vector normal to the boundary, $\mathbf{n} \cdot \mathbf{J}(t)$ is the net flux which traverses the boundary surface and A is the area of that boundary surface. We suggestively write the net flux $\mathbf{n} \cdot \mathbf{J}(t)$ in terms of $J_2^{in}(t)$ and $J_2^{out}(t)$, the probability fluxes in and out of the diffusion volume in question.

We see from comparison to (2) that (15) can be written out explicitly for our case

$$\{J_2^{in}(t) - J_2^{out}(t)\}A = k_{1 \rightarrow 2}P_1(t) - k_{2 \rightarrow 1}P_2(t). \quad (16)$$

One can easily see that the positive terms on either side of (16) are equal to each other as are the negative terms. Referring to (12), this quantity is

$$J_2^{out}(t)A = k_{2 \rightarrow 1}P_2(t) \quad (17)$$

making the rate

$$k_{2 \rightarrow 1} = \frac{J_2^{out}(t)A}{P_2(t)}. \quad (18)$$

In this expression the one-dimensional “area” through which the probability is flowing is simply a point and therefore $A = 1$ so identifying $J_2^{out}(t)$ with (13) we can solve for the unfolding rates in $\hat{K}$, $k_{2 \rightarrow 1}$ in (2) and (3), by dividing (13) by $P_2(t)$, or setting $P_2(t) = 1$ in (18) according to the stationary flux method. We find the backward folding rate in the one-dimensional case to be

$$k_{2 \rightarrow 1} = \frac{1}{L} \left(\frac{k_B T}{2\pi\mu} \right)^{1/2} e^{-E_b/k_B T}. \quad (19)$$

The terms preceding the exponential correspond to our prediction for the Van’t Hoff-Arrhenius attempt rate ν in (5). As an example, the attempt rate found for a coalesced pair of 16-residue Regan-Degrado (Regan and Degrado 1988) helices with a combined hydrophobic area loss of $600\text{\AA}^2$ is $64 \times 10^9 s^{-1}$.

We believe that the most probable dissociation of a microdomain-microdomain pairing occurs via relative motion along a vector connecting the centers-of-mass of the two microdomains. This implies that typically the one dimensional case is the best way to view the dissociation event and therefore that one should use (19) to go about calculating the rate. The one dimensional case should also be sufficient if there is an “unzipping” of paired α -helices.

It is possible, however, that dissociation may also include a relative rolling motion or other motion perpendicular to the axis connecting the microdomain pairs. In this case one needs to repeat the calculation above with a few minor differences: The relative velocity distribution of the microdomains is still the one dimensional Maxwell-Boltzmann distribution because the degrees of freedom parallel to the surface through which the probability is flowing do not contribute to escape from the well, the probability in the bound region is assumed to be evenly distributed in a two-dimensional “volume”, namely $\rho(x, t) = P_2(t)/\pi L^2$ in the two-dimensional analogue of (12), and the flux goes through a two-dimensional “area” $A = 2\pi L$ in (18). This calculation

yields the result

$$k_{2 \rightarrow 1} = \frac{2}{L} \left(\frac{k_b T}{2\pi\mu} \right)^{1/2} e^{-E_b/k_B T}. \quad (20)$$

Due to the steric clashing of the side chains it seems rather unlikely that dissociation would include a relative sliding motion along the axes of the microdomains. For completeness, however, we include the three-dimensional result derivable from analogous considerations to the ones given above

$$k_{2 \rightarrow 1} = \frac{3}{L} \left(\frac{k_b T}{2\pi\mu} \right)^{1/2} e^{-E_b/k_B T}. \quad (21)$$

Although the three-dimensional result seems an unlikely candidate for protein unfolding it may be relevant in the context of molten globules.

This approach succeeds in removing the free parameter ν from the model, and allows us to find the backward rates from a simple energetic model based on diffusion in a potential with appropriate boundary conditions. Moreover our results for the one, two and three-dimensional unfolding rates have a $\sqrt{T/\mu}$ dependence that could be used to distinguish between this and other proposals for the mechanism of helix-helix dissociation.

The removal of the parameter ν is important when considering folding processes which do not involve the native state. In previous applications of the diffusion-collision model, the folding kinetics from a denatured or random coil state to the final native state were followed. In such a case, it is reasonable to set the parameter ν such that the native state achieves 90 or 95 percent of the probability, because we know that the final state is attained at the end of the folding process. In studying intermediate processes or more importantly, non-native intermediates (Beck et al, 2000), where the occupation probability may be completely unknown, such reasonable estimates of ν are not available. In this case, elimination of ν as a free parameter is crucial.

III. Concluding Remarks

We have presented a calculation for the helix-helix dissociation rate using a simplified potential surface, a square potential having a depth equal to the free energy savings of hydrophobic docking and width equal to the diameter of a water molecule. We have found the unfolding rates arising from thermal fluctuations out of this potential well to be in good agreement with currently accepted values of ν . The potential itself is due to hydrophobic forces, which to date are not well understood and for this reason the potential chosen was simple.

The initial motivation of this work was to eliminate the free parameter ν from the diffusion-collision model. In the context of this model our result should allow us to perform kinetics simulations than were not possible before, for instance time developments of non-native intermediates, in which the occupation probabilities are unknown, and reasonable estimates for ν are not possible.

The results presented here also predict a $\nu \propto \sqrt{T/\mu}$ dependence in all cases which can be distinguished experimentally from other proposals such as the covalent bond model where $\nu \propto k_B T/\hbar$ (Fersht, 1999). Another difference is the dependence of the unfolding rates on the states, not only through the hydrophobic area, but also through the reduced mass μ of the microdomains or groups of microdomains undergoing dissociation. This is markedly different from typical diffusion-collision model calculations where the attempt rate ν is assumed to be the same for all dissociation events within the protein.

Along the way we have re-examined the idea that couching the problem of association and dissociation of microdomain pairs via diffusion over a potential surface affords us a clear and simple picture of the protein folding process. Indeed, the rate equations of the diffusion-collision model can be derived from such a picture. Interestingly, the potential energy surface is an element of other models such as folding funnels (for an overview see Bryngelson et al 1995).

Admittedly, our approximation is rather rough. The thrust of future work

should therefore be in constructing more realistic potentials including activation energies of hydrophobic dockings.

Generalization to more complex proteins is straightforward because every interaction between helices or clusters of helices can be considered a two-state process similar to the one described above; generally, the folding or unfolding of a given protein involves several such processes. In this case the diffusion space for each possible pairing could be constructed, and the forward and backward rates for each transition can be found, as outlined above, to construct the transition rate matrix $\hat{K}$ in order to find the time evolution of the state vector $\mathbf{P}(t)$. This is a much more tenable proposition than directly solving the Smoluchowski equation (7) on such a complicated potential surface, although it could be done in principle.

Acknowledgments

We would like to thank Ken Olum, David Weaver, and Larry Ford for many useful discussions. We are further grateful to David Weaver for pointing out a mistake in our calculations and Ken Olum for helping us fix it.

References

- Arrhenius S.** 1889. *Z Phys Chem* *4*:226.
- Bashford D, Cohen FE, Karplus M, Kuntz ID, Weaver DL** 1988. Diffusion-collision model for the folding kinetics of myoglobin. *Proteins* *4*:211-227.
- Beck C, Siemens X, Weaver DL.** 2000. Non-Native Kinetic intermediates and the Diffusion-Collision model of Protein Folding. *In preparation*.
- Burton RE, Meyers JK, Oas TG.** 1998. Protein folding dynamics: quantitative comparison between theory and experiment. *Biochemistry* *37*:5337-5343.
- Bryngelson JN, Onuchic ND, Socci ND, Wolynes PG.** 1995. Funnels,

pathways, and the energy landscape of protein folding: a synthesis. *Proteins: Struct Func and Gen* 21:need pages

Chandler D. 1987. Introduction to Modern Statistical Mechanics. Oxford University Press.

Chothia C. 1974. Hydrophobic bonding and accessible surface area in proteins. *Nature* 248:338-339.

Dinner AR, Lazaridis T, Karplus M. 1999. Understanding β -hairpin formation. *Proc Nat Acad Sci USA* 96:9068-9073.

Fersht A. Structure and mechanism in protein science. 158-159. W.H. Freeman and Co. New York. 1999.

Hanggi P, Talkner P, Borkovec M. 1990. Reaction-rate theory - 50 years after Kramers. *Rev Mod Phys* 62:251-341.

Karplus M, Weaver DL. 1976. Protein folding dynamics. *Nature* 260:404-406.

Karplus M, Weaver DL. 1979. Diffusion-collision model for protein folding. *Biopolymers* 18:1421-1437.

Lacroix E, Viguera AR, Serrano L. 1998. Elucidating the folding problem of α -helices: Local motifs, long-range electrostatics, ionic strength dependence, and prediction of of NMR parameters. *J Mol Biol* 284:173-191.

Lee B, Richards FM. 1971. The interpretation of protein structure: Estimation of static accessibility. *J Mol Biol* 55:379-400.

Munoz V, Henry ER, Hofrichter J, Eaton WA. 1998. A statistical mechanical model for β -hairpin kinetics. *Proc Nat Acad Sci USA* 95:5872-5879.

Munoz V, Serrano L. 1994a. Elucidating the folding problem of helical peptides using empirical parameters. *Nature: Struct Biol* 1:399-409.

Munoz V, Serrano L. 1994b. Elucidating the folding problem of helical peptides using empirical parameters II: Helix macrodipole effects and rational modification of the helical content of natural peptides. *J Mol Biol* 245:275-296.

Munoz V, Serrano L. 1994c. Elucidating the folding problem of helical

peptides using empirical parameters III: Temperature and pH dependence. *J Mol Biol* 245:297-308.

Munoz V, Serrano L. 1997. Development of the multiple sequence approximation within the AGADIR model of α -helix formation. Comparison with Zimm-Bragg and Lifson-Roig formalisms. *Biopolymers* 41:495-509.

Nagi AD, Anderson KS, Regan L 1999. Using loop length variant to dissect the folding pathway of a four-helix-bundle protein. *J Mol Biol* 286:257-265.

Pappu RV, Weaver DL. 1997. The early folding kinetics of apomyoglobin. *Protein Science* 7:480-490.

Regan L, Degrado WF. 1988. Characterization of a helical protein designed from first principles. *Science* 241:976-978.

Szabo A, Schulten K, Schulten Z. 1980. First passage time approach to diffusion controlled reactions. *J Chem Phys* 72:4350.

Van't Hoff JH. 1884. Studies in chemical dynamics. London, 1896.

Weaver DL. 1979. Some exact results for one-dimensional diffusion with absorption. *Phys Rev B* 20:2558.

Weiss GH. 1967. First passage time problems in chemical physics. *Adv Chem Phys* 13:1-18.

Zimm BH, Bragg JK. 1959. Theory of phase transition between helix and random coil in polypeptide chains. *J Chem Phys* 31:526-535.